

[Microbiol. Immunol., 38, 983-988 (1994)]

[Lab. of Pharmacology]

**Mechanism for macrophage activation against *Corynebacterium parvum*:  
participation of T cells and its lymphokines.**

HIROSHI MORI, MASAHIKO MIHARA, YUKI UESUGI, HIROICHI NAGAI\*, AKIHIDE KODA

Participation of T cells in the activation of macrophages by *C. parvum* was examined. TNF production in vitro from the spleen cells of BALB/c-+/+ mice was abrogated completely by the pre-treatment of spleen cells with anti-Ia antiserum and complement. TNF production was not elicited at all in BALB/c-nu/nucle mice. Supernatant from a culture of spleen cells stimulated with phytohemagglutinin-P (a PHA-induced lymphokine) made it possible for BALB/c-nu/nucle mice to produce TNF, associated with an induction of Lyt-1<sup>+</sup> cells and Lyt-2<sup>+</sup> cells. These results suggest that increasing the number of Ia<sup>+</sup> cells is not sufficient to bring about TNF production, and that Ia<sup>+</sup> cells must also be stimulated by T cells or T-cell lymphokines in order to produce TNF.

[Biol. Pharm. Bull., 17, 921-926 (1994)]

[Lab. of Pharmacology]

**Lymphocyte stimulation test with tetrazolium-based colorimetric assay for  
diagnosis of drug-induced allergic hepatitis.**

TOMOAKI YOSHIMURA, CHIKAKO KURITA, FUTOSHI YAMAZAKI, SATOSHI NAKANO,  
HIROICHI NAGAI\*

We examined the usefulness of 3-(4,5-dimethyl thiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) as a marker for the lymphocyte stimulation test (LST) in diagnosing drug allergy. A clinical study was made of 133 patients with diagnosed drug-induced allergic hepatitis. Among the causative drugs, antimicrobial agents were the most numerous accounting for 33.9 % of the total, followed by central nervous system agents 21.2 %, and cardiovascular agents 16.9 %. The MTT assay would be useful in diagnosing drug-induced allergic hepatitis, and among the drugs examined, antimicrobial agents were responsible for the largest number of allergic reactions.

[Planta Med., 60, 445-449 (1994)]

[Lab. of Pharmacology]

**Principle of the bark of *Phellodendron amurense* to suppress the cellular  
immune response.**

HIROSHI MORI, MASAHIRO FUCHIGAMI, NAOKI INOUE, HIROICHI NAGAI\*, AKIHIDE  
KODA, ITSUO NISHIOKA

OB-1 and OB-5, quaternary base alkaloids known as magnoflorine and phellodendrine, respectively, were isolated from *Phellodendri Cortex* as biologically active principles to suppress local graft-versus-host (GvH) reactions in mice. They suppressed the local GvH reaction, when given i.p. to the host mice at 5-20 mg/kg for 8 consecutive days from the day of spleen cell transfer. They also suppressed picryl chloride-induced delayed type hypersensitivity when given i.p. to mice at 10 and 20 mg/kg for 5 consecutive days from the day of the sensitization. These results suggest that OB-1 and OB-5 suppress the induction phase of the cellular immune response.