

[J. Pharmacol. Exp. Ther., 276, 231-237 (1996)]

[Lab. of Molecular Biology]

**A Catechol Derivative (4-Methylcatechol) Accelerates the Recovery
from Experimental Acrylamide-Induced Neuropathy**

KAZUKO SAITA, TAKEKAZU OHI, YASUO HANAOKA, SHOEI FURUKAWA*,
YOSHIKO FURUKAWA, KYOZO HAYASHI, SHIGERU MATSUKURA

Acrylamide (ACR) monomer produces neuropathy of the dying-back type and 4-methylcatechol (4-MC) is a potent stimulator of endogenous NGF synthesis. In the present study, we investigated the efficacy of 4-MC in promoting recovery from experimental ACR neuropathy in rats. Twenty-two Sprague-Dawley rats were made neuropathic by ACR injections. They showed hindlimb paralysis, increment of landing foot spread distance and a statistically significant reduction in motor nerve conduction velocity. 4-MC-administered ACR neuropathy rats showed improvement and increase in NGF content in the sciatic nerves. These findings suggest that 4-MC can accelerate the recovery process clinically.

[Tetrahedron Lett., 37, 7399-7402 (1996)]

[Lab. of Molecular Biology]

**Erinacines E, F, and G, Stimulators of Nerve Growth Factor
(NGF)-Synthesis, from the Mycelia of *Herichium erinaceum***

HIROKAZU KAWAGISHI, ATSUSHI SHIMADA, SATOSHI HOSOKAWA, HIRONOBU MORI,
HIDEKI SAKAMOTO, YUKIO ISHIGURO, SHINICHI SAKEMI, JON BORDNER,
NAKAO KOJIMA, SHOEI FURUKAWA*

The structure of erinacines E, F and G from mycelia of *Herichium erinaceum* were determined by spectroscopic and/or X-ray analysis. Erinacines E and F exhibited potent stimulating activity against NGF synthesis by astroglial cells.

[Heterocyclic Commun. , 2, 51-54 (1996)]

[Lab. of Molecular Biology]

**Erinacine D, A Stimulator of NGF-Synthesis, from the Mycelia of
*Hricium Erinaceum***

HIROKAZU KAWAGISHI, ATSUSHI SHIMADA, KAZUE SHIMIZU, HIRONOBU MORI,
KENJI OKAMOTO, HIDEKI SAKAMOTO, SHOEI FURUKAWA*

A novel diterpenoid, erinacine D, was isolated from the cultured mycelia of *Herichium erinaceum*. The structure of the compound was determined by interpretation of the spectral data and chemical reaction. This compound showed stimulating activity of nerve growth factor (NGF)- synthesis.