

[*J. Chem. Soc., Chem. Commun.*, 839-840 (2001)]

[Lab. of Pharm. Chemistry]

**Novel Acylation of a Vinyl Group by the Reaction of an Aldehyde and a Vinylselenonium Ylide.**

Shin-ichi WATANABE, Tomokazu KUSUMOTO, Chikayo YOSHIDA and Tadashi KATAOKA \*

Vinylselenonium ylide, which was generated from (*Z*)-vinylselenonium salt with a base such as sodium or potassium hydride, reacted with aromatic aldehydes to produce the  $\alpha,\beta$ -unsaturated ketones, which were obtained in better yields from the aldehydes with an electron-withdrawing group than from those with an electron-donating group. Based on the results from reactions using (vinyl-*d*)selenonium salts on benz(aldehyde-*d*) the reaction mechanism was proposed.

[*J. Chem. Soc., Chem. Commun.*, 1958-1959 (2001)]

[Lab. of Pharm. Chemistry]

**Self-assisted Tandem Michael-aldol Reactions of  $\alpha,\beta$ -Unsaturated Ketones with Aldehydes.**Tadashi KATAOKA, \* Sayaka KINOSHITA, Hironori KINOSHITA, Masaru FUJITA,  
Tatsunori IWAMURA and Shin-ichi WATANABE

The tandem Michael-aldol reaction of 1-[2-(methylsulfanyl)phenyl]prop-2-en-1-one (**1**) or the seleno congener **4** with *p*-nitrobenzaldehyde in the presence of  $\text{BF}_3 \cdot \text{OEt}_2$  gave the Baylis-Hillman adduct **2** or **5** and onium salt **3** or **6**, respectively, and selenochromanone **7** from **4**. When the reaction mixture of **1** or **4** with *p*-nitrobenzaldehyde was worked up with 2 equivalents of triethylamine, **2** (75%) or **5** (64%) together with **7** (10%) was obtained.

[*Heteroatom Chem.*, **12**, 317-326 (2001)]

[Lab. of Pharm. Chemistry]

**Syntheses and Reactions of Cyclic *Se*-Alkoxy-*Se*-chloroselenuranes and Alkoxy-selenonium Salts.**Tadashi KATAOKA, \* Tatsunori IWAMURA, Hisayoshi TSUTSUI, Yasuharu KATO,  
Yoshihiro BANNO, Yasuko AOYAMA, and Hiroshi SHIMIZU

Several 1-chloro-2,1-oxaselenole selenuranes, 5-chloro-5,11-epoxy-6,11-dihydrodibenzo[*b,e*]selenepines and the corresponding selenonium salts were synthesized and the differences in structures between the selenuranes and the selenonium were studied by  $^1\text{H}$ - and  $^{77}\text{Se}$ -NMR spectroscopy. Their reactions with organometallic reagents were conducted and the reaction mechanism was discussed.

[*Heterocycles*, **55**, 465-468 (2001)]

[Lab. of Pharm. Chemistry]

**Novel Ring Transformation of Dihydroselenines to Selenabicyclo[3.1.0]hexenes.**

Eiji HONDA, Shin-ichi WATANABE, Tatsunori IWAMURA and Tadashi KATAOKA \*

Treatment of 2-bromo-2,6-dicyano-2,3-dihydroselenine with triethylamine in ethanol gave 2-selenabicyclo[3.1.0]hex-3-ene (**1**) in 77% yield. Reaction of **1** with benzyne formed 2-cyanobenzoselenophene in 35% yield. Methylation of **1** with methyl triflate produced *Se*-methylselenonium salt (**2**), which was transformed into 2-selenabicyclo[3.1.0]hex-3-ene-1-carboxamide (**3**) and 3-carboxamide (**4**). Compound (**1**) was converted into cyanoethynylcyclopropane via selenonium salt (**2**).