

[Comp. Biochem. Physiol., 82B, 269 (1985)]

Characterization of Fe²⁺-Activated Acid Phosphatase in Rat Epidermis. AKIRA HARA, TAKASHI KATO, HIDEO SAWADA*, KIMIE FUKUYAMA, W. L. EPSTEIN

A particulate acid phosphatase (EC 3.1.3.2, orthophosphoric monoester phosphohydrolase (acid optimum)) was extracted in 1 M KCl, from 2-day-old rat epidermis. The enzyme has a Mr of 32,000, but two forms, F1 and F2 with pI values of 8.6 and 8.3, respectively, were identified while the pI values of other acid phosphatases soluble in sucrose and Triton X-100 were all acidic. F1 and F2 also differed from other epidermal acid phosphatases because they were (a) activated by Fe²⁺ and reducing agents, (b) showed immunological cross-reactivity with purple acid phosphatase of rat spleen and (c) dephosphorylated phosvitin and α -casein even though they had rather high Km values.

[Biochem. J., 225, 177 (1985)]

Guinea-Pig Liver Testosterone 17 β -Dehydrogenase (NADP⁺) and Aldehyde Reductase Exhibit Benzene Dihydrodiol Dehydrogenase Activity.

AKIRA HARA, MASAKAZU HAYASHIBARA, TOSHIHIRO NAKAYAMA, KAZUHISA HASEBE, SHIGEYUKI USUI, HIDEO SAWADA*

We have kinetically and immunologically demonstrated that testosterone 17 β -dehydrogenase isozymes and aldehyde reductase from guinea-pig liver catalyse the oxidation of benzene dihydrodiol to catechol. One isozyme of testosterone 17 β -dehydrogenase, which has specificity for 5 β -androstanes, oxidized benzene dihydrodiol at a 3-fold higher rate than 5 β -dihydrotestosterone, and showed a more than 4-fold higher affinity for benzene dihydrodiol and V_{max} value than did another isozyme, which exhibits specificity for 5 α -androstanes, and aldehyde reductase.

[J. Biochem., 98, 1131 (1985)]

Specificity of Hydrogen Transfer of Mammalian and Avian Carbonyl and Aldehyde Reductases. TOSHIHIRO NAKAYAMA, AKIRA HARA,

KIN-YA KARIYA, HIDEO SAWADA*

Aldehyde reductases from several mammalian tissues and chicken kidney transferred the *pro-4R* hydrogen of NADPH to the substrate, but the stereospecificity of carbonyl reductase from the tissues was not uniform. Carbonyl reductases from rat and guinea pig liver, which were associated with 3 α - and 17 β -hydroxysteroid dehydrogenase activities, were A-specific, whereas the enzymes from livers of man and monkey, guinea pig lung and chicken kidney, which did not exhibit the hydroxysteroid dehydrogenase activities, were all B-specific.