

[Heterocycles, 24, 2293 (1986)]

**Pyrimidines. Part 57. A Versatile Synthesis of Pyrimido [4,5-*d*]-  
pyrimidine-2,4,5-trione Derivatives**

KOSAKU HIROTA\*, JIAZI HUANG, HIRONAO SAJIKI, YOSHIFUMI MAKI

Bicyclic fused pyrimidines have been received much attention in connection with the biologically significant system such as purines and pteridines. Synthesis of the title compounds was accomplished by cyclization of 6-amino-1,3-dimethyl-5-(*N*-substituted carbamoyl or thiocarbamoyl) uracil derivatives (1), easily prepared from 6-amino-1,3-dimethyluracil and isocyanates, with one-carbon reagents. The treatment of (1) with dimethylformamide dimethylacetal, acetic anhydride, benzoic anhydride, and *N*, *N'*-carbonyldiimidazole gave the corresponding 7-*H*, methyl, phenyl, and hydroxypyrimido [4,5-*d*]-pyrimidines, respectively, in high yields.

[Tetrahedron Lett., 27, 3263 (1986)]

**Novel Reaction of 5-Nitro (or Carbamoyl) uracil Derivatives with Amines.  
Thermal Exchange Reaction of *N*<sub>1</sub>-position of the Uracils for Amines  
*via* Ring-Opening and Ring-Closure Processes**

KOSAKU HIROTA\*, YUKIO KITADE, HIRONAO SAJIKI, YOSHIFUMI MAKI

A novel type of reaction of uracils with amines under thermal conditions is described. The reaction of 1,3-disubstituted uracils (1) possessing an electron withdrawing group, such as nitro and carbamoyl groups, at the 5-position with ammonia and primary amines induced exchange of *N*<sub>1</sub>-portion of the uracil ring for amines. This reaction was accelerated by substitution of phenyl groups at the *N*<sub>1</sub>-position. The reaction intermediate, an amine-adduct of (1), was isolated. Two plausible reaction mechanisms for the exchange reaction are proposed on the basis of reactivity of the adduct.

[Chem. Pharm. Bull., 34, 664 (1986)]

**Reactions of 10-Thiaisoalloxazines with Alkyl Amines and Benzyl Alcohol.**

MAGOICHI SAKO, TAKASHI NIWA, KOSAKU HIROTA, YOSHIFUMI MAKI\*

Upon treatment with benzylamine and *n*-propylamine, 10-thiaisoalloxazines (1) undergo a smooth ring-contraction initiated by nucleophilic addition of alkyl amines to the conjugated diimine bond in the molecule to give the corresponding 6'-substituted amino-spiro [benzothiazoline-2,5'-pyrimidine-2',4' (3'*H*, 5'*H*)-diones], whereas thermal reaction of 1 with benzyl alcohol results in the preferential formation of the redox products, 1,5-dihydro-10-thiaisoalloxazines and benzaldehyde, proceeding via an initial single-electron transfer process. The latter result is suggestive of the presence of an alternative process not involving the carbanion intermediates in the flavin-catalyzed oxidation of alcohols.