

[J. Biochem., 106, 900 (1989)]

3(20) α -Hydroxysteroid Dehydrogenase Activity of Monkey Liver Indanol Dehydrogenase.

AKIRA HARA, MAKOTO NAKAGAWA, HIROYUKI TANIGUCHI,
HIDEO SAWADA*

Homogeneous indanol dehydrogenase from monkey liver oxidized reversibly 3 α - or 20 α -hydroxy groups of several bile acids and 5 β -pregnanes with high k_{cat}/K_m values. Competitive inhibition of the hydroxysteroid dehydrogenase activity of the enzyme by medroxyprogesterone acetate, hexestrol, and 1,10-phenanthroline suggests that both 1-indanol and hydroxysteroid are oxidized at the same active site on the enzyme. The specific inhibitor of the enzyme, 1,10-phenanthroline, suppressed the 3 α -hydroxysteroid dehydrogenase activity in the crude extract of monkey liver by 50%. The results strongly suggest that indanol dehydrogenase acts as a 3(20) α -hydroxysteroid dehydrogenase in the metabolism of certain steroid hormones and bile acids.

[J. Biochem., 106, 1104 (1989)]

Dimeric Dihydrodiol Dehydrogenase in Monkey Kidney. Substrate Specificity, Stereospecificity of Hydrogen Transfer, and Distribution.

MAKOTO NAKAGAWA, KAZUYA MATSUURA, AKIRA HARA, HIDEO SAWADA,*
YASUO BUNAI, ISAO OHYA

Chemical cross-linked and NADPH binding studies suggested that the native dihydrodiol dehydrogenase from monkey kidney is a basic dimer (Mr 78,000) and one active site per the subunit. The enzyme oxidized specifically *trans*-dihydrodiols of benzene and naphthalene, whereas it catalyzed the reduction of dihydroxyacetone phosphate at pH 7.4. The enzyme transferred the 4-pro-*R* hydrogen atom of NADPH to carbonyl substrate. Immunochemical and immunohistochemical studies using a specific antibody revealed that this enzyme specifically distributed in proximal and distal tubules of the cortex, and in the loop of Henle of the medulla in the kidney.

[Biochem. J., 264, 403 (1989)]

Isolation from Pig Lens of Two Proteins with Dihydrodiol Dehydrogenase and Aldehyde Reductase Activities.

AKIRA HARA, TATSUHIRO HARADA, MAKOTO NAKAGAWA,
KAZUYA MATSUURA, TOSHIHIRO NAKAYAMA, HIDEO SAWADA*

Dimeric and monomeric proteins containing dihydrodiol dehydrogenase and aldehyde reductase activities were purified from pig lens. The dimeric enzyme of Mr 65,000 specifically oxidized the *trans*-dihydrodiols of naphthalene and benzene with NADP⁺ as a strict cofactor, and reduced α -diketones, aromatic aldehydes and glyceraldehyde with NADPH as a cofactor. The monomeric enzyme of Mr 35,000, although identical with aldose reductase, oxidized the *trans*-dihydrodiol of naphthalene at a pH optimum of 7.6. These results suggest that the two enzymes are involved in the pathogenesis of naphthalene cataract.