

[Tetrahedron, 46, 7393 (1990)]

Hard Acid and Soft Nucleophile Systems. Part 12. Regioselective Functionalization of 1,3-Dienes through the Lewis Acid Mediated Thienium Cation Diels-Alder Reaction.

KAORU FUJI, SUBHASH P. KHANAPURE, MANABU NODE, TAKEO KAWABATA,
AKICHIKA ITOH*, YUKIO MASAKI

In the Diels-Alder reaction, the electron-donating alkyl sulfenyl groups at the β -position of α -nitroalkenes may decrease the normal reactivity toward dienes. However, we found that the use of aluminium chloride in the attempted Diels-Alder reaction of β -ethylthio- α -nitroalkenes with 1,3-dienes dramatically changed the reaction course. The reaction led to the formation of *Z*-olefins by the regio- and stereoselective 1,4-functionalization of 1,3-dienes, which probably involve the Diels-Alder cycloaddition of a thienium cation intermediates, generated from the β -ethylthio- α -nitroalkenes, with the dienes.

[J. Chem. Soc., Faraday Trans., 86, 2971 (1990)]

Physicochemical Properties of a Novel Nitrogen-containing Plasma-deposited Ultrathin Film.

MASAYUKI KUZUYA*, AKIHIRO NOGUCHI, YOSHIKAZUMI TANAKA,
SACHIIHIKO NAKAI, KAORU KAMIYA, YASUKICHI YANAGIHARA

Amphiphilic nitrogen-containing ultrathin films have been formed by brief plasmolysis of various kinds of nitrogen-containing organic compounds such as amino-, amido-, nitrile- and nitro-derivatives. Ultrathin films prepared from nitrogen-rich organic compounds, however, have been found to induce neither effective polymerization of vinyl monomers nor spin adduct formation of PBN, unlike those from non-nitrogen-containing organic compounds. Most of these films were very unstable in solvents, and the nitrogen-containing moiety in the resulting films was readily leached out of the film into either polar or non-polar solvents.

[J. Chem. Soc., Faraday Trans., 86, 2975 (1990)]

Effects and Role of Basic Nitrogen-derived Ultrathin Films on 'Plasma-Reduction'.

MASAYUKI KUZUYA*, AKIHIRO NOGUCHI, SEIKO MIZUTANI,
YOSHIKAZUMI TANAKA

Basic nitrogen-derived ultrathin films have been formed on brief plasmolysis (in a closed system) of even non-basic nitrogen-containing organic vapours such as dimethylformamide (DMF) and nitrile derivatives. The nitrogen-containing entity in these is readily leached into either polar or non-polar solvents. Based on these findings, the reported 'plasma reduction' of various electron-accepting compounds by energetic electrons could be rationalized in terms of nitrogen base-catalysed reactions, since various viologens, organic dye compounds, iodine and all other compounds reported for 'plasma reduction' also undergo electronic spectral changes in reactions with nitrogen bases.